

For adults with moderate to severe submental fat

# DIMINISH THE DOUBLE CHIN

*Real patient. Results may vary.*

Reduce moderate to severe fullness under your chin and reveal an improved chin profile.

### What is KYBELLA®?

KYBELLA® is a prescription medicine used in adults to improve the appearance and profile of moderate to severe fat below the chin (submental fat), also called “double chin.”

It is not known if KYBELLA® is safe and effective for the treatment of fat outside of the submental area or in children under 18 years of age.

### Important Safety Information Who should not receive KYBELLA®?

Do not receive KYBELLA® if you have an infection in the treatment area.



**Please see additional Important Safety Information throughout this document.  
View full Prescribing Information or visit [https://www.rxabbvie.com/pdf/kybella\\_pi.pdf](https://www.rxabbvie.com/pdf/kybella_pi.pdf).**

 **kybella**<sup>®</sup>  
(deoxycholic acid) injection 10 mg/mL

**First-of-its-kind nonsurgical treatment** for moderate to severe fullness under the chin

**Permanently destroys fat cells** in the treatment area

A series of **tailored treatments**\*

\*Multiple injections under the chin per treatment; up to 6 treatments spaced at least 1 month apart. 59% of patients received 6 KYBELLA<sup>®</sup> treatments in clinical studies.

Is KYBELLA<sup>®</sup> right for you?  
Ask your specialist today.

*Real patient. Results may vary.*

## WHAT TO EXPECT With KYBELLA<sup>®</sup>

- **The area under and around your chin will be examined** to determine if KYBELLA<sup>®</sup> is right for you. Your specialist may ask some questions about your medical history
- **You'll discuss your aesthetic goals** with your specialist and **receive a tailored treatment plan**. Expect between 2 and 6 treatment sessions, spaced at least 1 month apart\*
- At each session, you'll receive multiple small injections under your chin. **The most common side effects include swelling, pain, numbness, redness, and areas of hardness in the treatment area**. Return for all subsequent treatments to help achieve your goals

### Important Safety Information (continued)

**Before receiving KYBELLA<sup>®</sup>, tell your healthcare provider about all of your medical conditions, including if you:** Have had or plan to have surgery on your face, neck, or chin; have had cosmetic treatments on your face, neck, or chin; have had or have medical conditions in or near the neck area; have had or have trouble swallowing; have bleeding problems; are pregnant or plan to become pregnant (it is not known if KYBELLA<sup>®</sup> will harm your unborn baby); are breastfeeding or plan to breastfeed (it is not known if KYBELLA<sup>®</sup> passes into your breast milk).

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take a medicine that prevents the clotting of your blood (antiplatelet or anticoagulant medicine).

**Please see additional Important Safety Information throughout this document. View full Prescribing Information or visit [https://www.rxabbvie.com/pdf/kybella\\_pi.pdf](https://www.rxabbvie.com/pdf/kybella_pi.pdf).**

 **kybella**<sup>®</sup>  
(deoxycholic acid) injection 10 mg/mL

# ARE YOU A 2+ OR 4+ CANDIDATE?

## 2+

TREATMENTS

Expect 2 or 3 treatments to achieve desired aesthetic goals

ADRA  
AGE: 35

BEFORE

3 TREATMENTS

MARINA  
AGE: 58

BEFORE

3 TREATMENTS

## 4+

TREATMENTS

Expect 4, 5, or 6 treatments to achieve desired aesthetic goals

PONTI  
AGE: 48

BEFORE

6 TREATMENTS

JASON  
AGE: 36

BEFORE

6 TREATMENTS

Unretouched photos of paid models. Individual results may vary. Not all treatments are shown; 59% of adults received 6 KYBELLA® treatments in clinical studies.

### Important Safety Information (continued)

#### What are the possible side effects of KYBELLA?


**KYBELLA® can cause serious side effects, including**

- Nerve injury in the jaw (which can cause an uneven smile or facial muscle weakness)
- Trouble swallowing
- Injection site problems including: a collection of blood under the skin (hematoma) or bruising, damage to an artery or vein if KYBELLA® is inadvertently injected into it, hair loss, open sores (ulcers), damage and tissue cell-death (necrosis) around the injection site. Call your healthcare provider if you: begin to develop weakness in the muscles of your face, or your smile becomes uneven; have difficulty swallowing, or if any of the symptoms that you already have get worse; develop open sores or drainage from the treatment area

Please see additional Important Safety Information throughout this document. View full Prescribing Information or visit [https://www.rxabbvie.com/pdf/kybella\\_pi.pdf](https://www.rxabbvie.com/pdf/kybella_pi.pdf).

## Tailor your treatment plan.

Talk to your specialist about the number of treatments you may need.

 **kybella**<sup>®</sup>  
(deoxycholic acid) injection 10 mg/mL

 **kybella**<sup>®</sup>  
(deoxycholic acid) injection 10 mg/mL

For adults with moderate to severe submental fat

**DIMINISH THE  
DOUBLE CHIN**

**Allē**

*Earn on the Things You Love*

Allē rewards you with points on Allergan Aesthetics brands and a variety of in-office treatments from other brands too.

Visit [Alle.com](https://alle.com)

**Important Safety Information (continued)**

**The most common side effects of KYBELLA<sup>®</sup> include** swelling, pain, numbness, redness, and areas of hardness in the treatment area.

These are not all of the possible side effects of KYBELLA<sup>®</sup>. Call your doctor for medical advice about side effects.

Ask your healthcare provider or visit [MyKybella.com](https://MyKybella.com) for full Prescribing Information.

**Allergan  
Aesthetics**  
an AbbVie company

© 2022 AbbVie. All rights reserved. KYBELLA<sup>®</sup> and its design are registered trademarks of Allergan Sales, LLC, an AbbVie company. KYB153025 01/22 014909

 **kybella**<sup>®</sup>  
(deoxycholic acid) injection 10 mg/mL

*Real patient. Results may vary.*